

SPECIALTY GUIDELINE MANAGEMENT

CIMZIA (certolizumab pegol)

POLICY

I. INDICATIONS

The indications below including FDA-approved indications and compendial uses are considered a covered benefit provided that all the approval criteria are met and the member has no exclusions to the prescribed therapy.

A. FDA-Approved Indications

1. Moderately to severely active rheumatoid arthritis (RA)
2. Active psoriatic arthritis (PsA)
3. Active ankylosing spondylitis (AS)
4. Moderately to severely active Crohn's disease (CD)
5. Moderate to severe plaque psoriasis (PsO)

B. Compendial Uses

Axial spondyloarthritis

All other indications are considered experimental/investigational and therefore considered not medically necessary.

II. CRITERIA FOR INITIAL APPROVAL

A. **Moderately to severely active rheumatoid arthritis (RA)**

1. Authorization of 12 months may be granted for members who have previously received Cimzia or any other biologic DMARD or targeted synthetic DMARD indicated for moderately to severely active rheumatoid arthritis and documentation confirms treatment failure or contraindication to Remicade and Simponi Aria.
2. Authorization of 12 months may be granted for treatment of moderately to severely active RA when documentation confirms treatment failure or contraindication to Remicade or Simponi Aria and any of the following criteria is met:
 - a. Member has experienced an inadequate response to at least a 3-month trial of methotrexate despite adequate dosing (i.e., titrated to 20 mg/week).
 - b. Member has an intolerance or contraindication to methotrexate (see Appendix A).

B. **Active psoriatic arthritis (PsA)**

Authorization of 12 months may be granted for treatment of active psoriatic arthritis (PsA) when documentation confirms treatment failure or contraindication to Remicade or Simponi Aria.

C. **Active ankylosing spondylitis (AS) and axial spondyloarthritis**

1. Authorization of 12 months may be granted for members who have previously received Cimzia or any other biologic DMARD indicated for active ankylosing spondylitis and documentation confirms treatment failure or contraindication to Remicade or Simponi Aria.

Reference number(s)
2005-A

2. Authorization of 12 months may be granted for treatment of active ankylosing spondylitis and axial spondyloarthritis when any of the following criteria is met:
 - a. Member has experienced an inadequate response to at least two non-steroidal anti-inflammatory drugs (NSAIDs).
 - b. Member has an intolerance or contraindication to two or more NSAIDs.

D. Moderately to severely active Crohn's disease (CD)

1. Authorization of 12 months may be granted for members who have previously received Cimzia or any other biologic indicated for the treatment of Crohn's disease and documentation confirms treatment failure or contraindication to Remicade.
2. Authorization of 12 months may be granted for treatment of moderately to severely active CD when documentation confirms treatment failure or contraindication to Remicade and the member has an inadequate response, intolerance or contraindication to at least one conventional therapy option (see Appendix B).

E. Moderate to severe plaque psoriasis (PsO)

1. Authorization of 12 months may be granted for members who have previously received Cimzia, or any other biologic DMARD indicated for the treatment of moderate to severe chronic plaque psoriasis and documentation confirms treatment failure or contraindication to Remicade.
2. Authorization of 12 months may be granted for treatment of moderate to severe chronic plaque psoriasis when all of the following criteria are met:
 - a. At least 5% of body surface area (BSA) is affected OR crucial body areas (e.g., hands, feet, face, neck, scalp, genitals/groin, intertriginous areas) are affected.
 - b. Member meets any of the following criteria:
 - i. Member has had an inadequate response or intolerance to either phototherapy (e.g., UVB, PUVA) or a pharmacologic treatment with methotrexate, cyclosporine or acitretin.
 - ii. Member has a clinical reason to avoid pharmacologic treatment with methotrexate, cyclosporine or acitretin (see Appendix C).
 - iii. Member has severe psoriasis that warrants a biologic DMARD as first-line therapy.

III. CONTINUATION OF THERAPY

Authorization of 12 months may be granted for all members (including new members) who meet all initial authorization criteria and achieve or maintain positive clinical response after at least 3 months of therapy with Cimzia as evidenced by low disease activity or improvement in signs and symptoms of the condition.

IV. OTHER

For all indications: Member has a pretreatment tuberculosis (TB) screening with a TB skin test or an interferon gamma release assay (e.g., QFT-GIT, T-SPOT.TB).

Note: Members who have received Cimzia or any other biologic DMARD or targeted synthetic DMARD are exempt from requirements related to TB screening in this Policy.

V. APPENDICES

Appendix A: Examples of Contraindications to Methotrexate

1. Alcoholism, alcoholic liver disease or other chronic liver disease
2. Breastfeeding
3. Blood dyscrasias (e.g., thrombocytopenia, leukopenia, significant anemia)
4. Elevated liver transaminases
5. History of intolerance or adverse event
6. Hypersensitivity
7. Interstitial pneumonitis or clinically significant pulmonary fibrosis
8. Myelodysplasia
9. Pregnancy or planning pregnancy (male or female)
10. Renal impairment
11. Significant drug interaction

Appendix B: Examples of Conventional Therapy Options for CD

1. Mild to moderate disease – induction of remission:
 - a. Oral budesonide
 - b. Alternatives: metronidazole, ciprofloxacin, rifaximin
2. Mild to moderate disease – maintenance of remission:
 - a. Azathioprine, mercaptopurine
 - b. Alternatives: oral budesonide, methotrexate intramuscularly (IM) or subcutaneously (SC), sulfasalazine
3. Moderate to severe disease – induction of remission:
 - a. Prednisone, methylprednisolone intravenously (IV)
 - b. Alternatives: methotrexate IM or SC
4. Moderate to severe disease – maintenance of remission:
 - a. Azathioprine, mercaptopurine
 - b. Alternative: methotrexate IM or SC
5. Perianal and fistulizing disease – induction of remission:
 - a. Metronidazole ± ciprofloxacin, tacrolimus
6. Perianal and fistulizing disease – maintenance of remission:
 - a. Azathioprine, mercaptopurine
 - b. Alternative: methotrexate IM or SC

Appendix C: Examples of Clinical Reasons to Avoid Pharmacologic Treatment with Methotrexate, Cyclosporine or Acitretin.

1. Alcoholism, alcoholic liver disease or other chronic liver disease
2. Breastfeeding
3. Drug interaction
4. Cannot be used due to risk of treatment-related toxicity
5. Pregnancy or planning pregnancy (male or female)
6. Significant comorbidity prohibits use of systemic agents (examples include liver or kidney disease, blood dyscrasias, uncontrolled hypertension)

VI. REFERENCES

1. Cimzia [package insert]. Smyrna, GA: UCB, Inc.; June 2018.
2. van der Heijde D, Ramiro S, Landewe R, et al. 2016 Update of the international ASAS-EULAR management recommendations for axial spondyloarthritis. *Ann Rheum Dis.* 2017;0:1-14.

Reference number(s)
2005-A

3. Smolen JS, Landewé R, Billsma J, et al. EULAR recommendations for the management of rheumatoid arthritis with synthetic and biological disease-modifying antirheumatic drugs: 2016 update. *Ann Rheum Dis.* 2017;0:1-18.
4. Singh JA, Saag KG, Bridges SL Jr, et al. 2015 American College of Rheumatology Guideline for the Treatment of Rheumatoid Arthritis. *Arthritis Rheumatol.* 2016;68(1)1-26.
5. Saag KG, Teng GG, Patkar NM, et al. American College of Rheumatology 2008 recommendations for the use of nonbiologic and biologic disease-modifying antirheumatic drugs in rheumatoid arthritis. *Arthritis Rheum.* 2008;59(6):762-784.
6. Menter A, Korman NJ, Elmets CA, et al. Guidelines of care for the management of psoriasis and psoriatic arthritis. Section 6: Guidelines of care for the treatment of psoriasis and psoriatic arthritis: case-based presentations and evidence-based conclusions. *J Am Acad Dermatol.* 2011;65(1):137-174.
7. Gossec L, Smolen JS, Ramiro S, et al. European League Against Rheumatism (EULAR) recommendations for the management of psoriatic arthritis with pharmacological therapies; 2015 update. *Ann Rheum Dis.* 2016;75(3):499-510.
8. Gladman DD, Antoni C, P Mease, et al. Psoriatic arthritis: epidemiology, clinical features, course, and outcome. *Ann Rheum Dis.* 2005;64(Suppl II):ii14–ii17.
9. Peluso R, Lervolino S, Vitiello M, et al. Extra-articular manifestations in psoriatic arthritis patients. [Published online ahead of print May 8, 2014]. *Clin Rheumatol.* 2014.
10. Braun J, van den Berg R, Baraliakos X, et al. 2010 update of the ASAS/EULAR recommendations for the management of ankylosing spondylitis. *Ann Rheum Dis.* 2011;70:896–904.
11. Landewe R, Braun J, Deodhar A, et al. Efficacy of certolizumab pegol on signs and symptoms of axial spondyloarthritis including ankylosing spondylitis: 24-week results of a double-blind randomised placebo-controlled Phase 3 study. *Ann Rheum Dis.* 2014;73(1):39-47.
12. Ward MM, Deodhar A, Akl EA, et al. American College of Rheumatology/Spondylitis Association of America/Spondyloarthritis Research and Treatment Network 2015 recommendations for the treatment of ankylosing spondylitis and nonradiographic axial spondyloarthritis. *Arthritis Rheumatol.* 2015: 10.1002/art.39298. [Epub ahead of print].
13. Talley NJ, Abreu MT, Achkar J, et al. An evidence-based systematic review on medical therapies for inflammatory bowel disease. *Am J Gastroenterol.* 2011;106(Suppl 1):S2-S25.
14. Lichtenstein GR, Loftus Jr EV, Isaacs KI, et al. ACG Clinical Guideline: Management of Crohn's Disease in Adults. *Am J Gastroenterol.* 2018;113:481-517.