

Harvard Pilgrim Health Care 2018 HEDIS Results: HMO/POS Commercial

(Calendar year 2017 performance – Reported in October 2018)

Table shows percent of sample meeting HEDIS or CAHPS measurement criteria (see companion Measure Guide for details)

| | HPHC, Inc. | National |
|---|------------|----------|
| | (MA, ME) | Average |
| ALL ADULTS | | J |
| Adults' Access to preventive/ambulatory health services | | |
| Ages 20 – 44 | 96% | 92% |
| Ages 45 - 64 | 98% | 96% |
| Ages 65+ | 99% | 97% |
| Ages total | 97% | 94% |
| Adult BMI assessment | 81% | 80% |
| Avoidance of antibiotic treatment for adults with acute | | |
| bronchitis | 56% | 31% |
| Colorectal cancer screening | 73% | 62% |
| Disease modifying anti-rheumatic drug therapy in | 91% | 89% |
| Rheumatoid Arthritis | | |
| Use of imaging studies for low back pain | 84% | 75% |
| Use of spirometry testing in the assessment of COPD | 40% | 42% |
| Flu vaccinations for adults* (ages 18-64) | 58% | 49 % |
| Advising Smokers to Quit* (ASTQ) | ٨ | |
| WOMEN & TEENAGE GIRLS | | |
| Breast cancer screening (mammograms) | 84% | 73% |
| Cervical cancer screening (pap smears) | 88% | 76% |
| Chlamydia screening in women | 3370 | 7070 |
| Ages 16-20 | 72% | 45% |
| Ages 21-24 | 74% | 55% |
| Overall (Ages 16-24) | 73% | 50% |
| Prenatal care in the first trimester of pregnancy | 90% | 84% |
| Postpartum visit after delivery | 80% | 75% |
| CHILDREN & ADOLESCENTS | 3373 | 7.0 7.0 |
| Appropriate treatment for children with upper respiratory | 97% | 89% |
| infections | J 7 7 0 | 0570 |
| Appropriate testing for children with pharyngitis (sore throat) | 97% | 87% |
| Childhood Immunization Status | 7770 | 0770 |
| Combo 10 (DTaP, IPV, MMR, HIB, Hep B, VZV, PCV, Hep A, | 70% | 54% |
| Rotavirus, Influenza) | 7070 | 3.70 |
| Immunizations for Adolescents | | |
| Meningococcal | 90% | 82% |
| Tdap | 95% | 89% |
| HPV | 27% | 24% |
| Combo 1 (Meningococcal & Tdap) | 90% | 80% |
| Combo 2 (Meningococcal, Tdap & HPV) | 26% | 22% |
| Children and Adolescents' Access to primary care physicians | 2,72 | |
| 12 months – 24 months | 100% | 98% |
| 25 months – 6 years | 98% | 93% |

^{*} Measure from the CAHPS (Consumer Assessment of Healthcare Providers and Systems) survey, not HEDIS (Healthcare Effectiveness Data and Information Set)

[^] ASTQ: Did not hit minimum number of respondents to calculate two-year rolling average

[~] Denominator less than required & therefore rate not reportable



Harvard Pilgrim Health Care 2018 HEDIS Results: HMO/POS Commercial

(Calendar year 2017 performance – Reported in October 2018)

| (Catenaar year 2017 perjormance – Reportea in Ol | HPHC, Inc. (MA, ME) | National Average |
|--|------------------------|---------------------|
| 7 years – 11 years | 99% | 93% |
| 12 years – 19 years | 98% | 91% |
| Well-care visits | 7070 | <i>J</i> 170 |
| 1 st 15 months | 95% | 81% |
| 3 years – 6 years | 93% | 79% |
| 12 years –21 years | 78% | 50% |
| Weight assessment & counseling for children | 7070 | 3070 |
| BMI percentile (all ages) | 82% | 71% |
| Nutrition counseling (all ages) | 83% | 66% |
| Physical activity counseling (all ages) | 77% | 60% |
| CARDIAC CARE (Adults Only) | , | |
| Persistence of beta-blocker treatment after a heart attack | 88% | 85% |
| ADULTS WITH HYPERTENSION | | |
| Controlling high blood pressure | 71% | 60% |
| ADULTS WITH DIABETES | | |
| Comprehensive diabetes care | | |
| Hemoglobin A1c (HbAlc) screening conducted | 92% | 91% |
| HbAlc control <8.0 (higher is better) | 61% | 57% |
| Dilated Eye Exam conducted | 74% | 54% |
| Medical Attention for Nephropathy (kidney disease) | 92% | 90% |
| Blood Pressure control <140/90 | 70% | 61% |
| ASTHMA TREATMENT (Adults & Children) | | |
| Asthma Medication Ratio | | |
| Ages 5-11 | 90% | 87% |
| Ages 12-18 | 80% | 77% |
| Ages 19-50 | 70% | 74% |
| Ages 51-64 | 81% | 83% |
| All ages | | |
| | 77% | 79% |
| BEHAVIORAL HEALTH CARE (Adults only) | | |
| Antidepressant medication management | | |
| Effective acute-phase treatment | 69% | 67% |
| Effective continuation-phase treatment | 53% | 52% |
| Follow-up appt. after hospitalization for mental illness | | |
| Within seven days of discharge | 66% | 47% |
| Within 30 days of discharge | 85% | 67% |
| Initiation of alcohol and other drug treatment - all ages | 34% | 36% |
| Engagement of alcohol and other drug treatment - all ages | 11% | 13% |

^{*} Measure from the CAHPS (Consumer Assessment of Healthcare Providers and Systems) survey, not HEDIS (Healthcare Effectiveness Data and Information Set)

[^] ASTQ: Did not hit minimum number of respondents to calculate two-year rolling average

[~] Denominator less than required & therefore rate not reportable