

Harvard Pilgrim Insurance Company (HPIC)

2018 HEDIS Results: PPO Commercial

(Calendar year 2017 performance – Reported in October 2018)

%

Table shows percent of sample meeting HEDIS or CAHPS measurement criteria (see companion *Measure Guide* for details)

	HPIC (CT)	National Average
ALL ADULTS		
Adults' Access to preventive/ambulatory health services		
Ages 20 – 44	90%	92%
Ages 45 - 64	94%	96%
Ages 65+	~	96%
Ages total	93%	94%
Adult BMI assessment	66%	67%
Avoidance of antibiotic treatment for adults with acute bronchitis	23%	30%
Colorectal cancer screening	46%	59%
Disease modifying anti-rheumatic drug therapy in Rheumatoid Arthritis	85%	88%
Use of imaging studies for low back pain	73%	76%
Use of spirometry testing in the assessment of COPD	~	40%
Flu vaccinations for adults* (ages 18-64)	48%	50%
Advising Smokers to Quit* (ASTQ)	^	
WOMEN & TEENAGE GIRLS		
Breast cancer screening (mammograms)	70%	70%
Cervical cancer screening (pap smears)	74%	73%
Chlamydia screening in women		
Ages 16-20	53%	42%
Ages 21-24	64%	52%
Overall (Ages 16-24)	59%	47%
Prenatal care in the first trimester of pregnancy	90%	77%
Postpartum visit after delivery	89%	67%
CHILDREN & ADOLESCENTS		
Appropriate treatment for children with upper respiratory infections	91%	88%
Appropriate testing for children with pharyngitis (sore throat)	95%	86%
Childhood Immunization Status		
Combo 10 (DTaP, IPV, MMR, Hib, Hep B, VZV, PCV, Hep A, Rotavirus, Influenza)	62%	48%
Immunizations for Adolescents		
Meningococcal	84%	75%
TDAP	90%	83%
HPV	14%	22%
Combo 1 (Meningococcal & TDAP)	84%	73%
Combo 2 (Meningococcal, TDAP & HPV)	12%	20%
Children and Adolescents' Access to primary care physicians		
12 months – 24 months	99%	97%

* Measure from the CAHPS (Consumer Assessment of Healthcare Providers and Systems) survey, not HEDIS (Healthcare Effectiveness Data and Information Set)

^ ASTQ: Did not hit minimum number of respondents to calculate two-year rolling average

~ Denominator less than required & therefore rate not reportable

Harvard Pilgrim Insurance Company (HPIC) 2018 HEDIS Results: PPO Commercial

(Calendar year 2017 performance – Reported in October 2018)

	HPIC (CT)	National Average
25 months – 6 years	97%	91%
7 years – 11 years	99%	91%
Children/Adolescents' Access (cont'd):		
12 years – 19 years	96%	89%
Well-care visits		
1 st 15 months	83%	78%
3 years – 6 years	87%	75%
12 years – 21 years	68%	45%
Weight assessment & counseling for children		
BMI percentile (all ages)	82%	57%
Nutrition counseling (all ages)	77%	53%
Physical activity counseling (all ages)	72%	48%
CARDIAC CARE (Adults Only)		
Persistence of beta-blocker treatment after a heart attack	~	84%
ADULTS WITH HYPERTENSION		
Controlling high blood pressure	50%	54%
ADULTS WITH DIABETES		
Comprehensive diabetes care		
Hemoglobin A1c (HbA1c) screening conducted	90%	90%
HbA1c control <8.0 (higher is better)	55%	48%
Dilated Eye Exam conducted	54%	49%
Medical Attention for Nephropathy (kidney disease)	90%	88%
Blood Pressure control <140/90	57%	50%
ASTHMA TREATMENT (Adults & Children)		
Asthma Medication Ratio		
Ages 5-11	~	88%
Ages 12-18	~	79%
Ages 19-50	68%	75%
Ages 51-64	82%	83%
Ages 65-85	NA	NA
All ages	79%	80%
BEHAVIORAL HEALTH CARE (Adults only)		
Antidepressant medication management		
Effective acute-phase treatment	74%	68%
Effective continuation-phase treatment	60%	53%
Follow-up appt. after hospitalization for mental illness		
Within seven days of discharge	67%	45%
Within 30 days of discharge	83%	67%
Initiation of alcohol and other drug treatment - all ages	37%	37%
Engagement of alcohol and other drug treatment - all ages	14%	14%

* Measure from the CAHPS (Consumer Assessment of Healthcare Providers and Systems) survey, not HEDIS (Healthcare Effectiveness Data and Information Set)

^ ASTQ: Did not hit minimum number of respondents to calculate two-year rolling average

~ Denominator less than required & therefore rate not reportable